


DISTANCES IN ORIENTATIONS OF GRAPHS

by

V. Chvatal  
G. Thomassen

STAN-CS-75-517  
AUGUST 1975

COMPUTER SCIENCE DEPARTMENT  
School of Humanities and Sciences  
STANFORD UNIVERSITY



# Distances in Orientations of Graphs

by

V. Chvátal and C. Thomassen

Computer Science Department  
Stanford University  
Stanford, California 94305

and

Department of Combinatorics and Optimization  
University of Waterloo  
Waterloo, Ontario, Canada

## Abstract

We prove that there is a function  $h(k)$  such that every undirected graph  $G$  admits an orientation  $H$  with the following property: if an edge  $uv$  belongs to a cycle of length  $k$  in  $G$  then  $uv$  or  $vu$  belongs to a directed cycle of length at most  $h(k)$  in  $H$ . Next, we show that every undirected bridgeless graph of radius  $r$  admits an orientation of radius at most  $r^2+r$ , and this bound is best possible. We consider the same problem with radius replaced by diameter. Finally, we show that the problem of deciding whether an undirected graph admits an orientation of diameter (resp. radius) two belongs to a class of problems called NP-hard.

This research was supported in part by National Science Foundation grant DCR72-03752 A02 and by the Office of Naval Research contract NR 044-402. Reproduction in whole or in part is permitted for any purpose of the United States Government.

0. Introduction.

In 1939, H. E. Robbins [12] proved that an undirected graph  $G$  admits a strongly connected orientation if and only if  $G$  is connected and bridgeless. If  $G$  is thought of as the system of two-way streets in a city, then the theorem gives necessary and sufficient conditions for being able to make every street in the city one-way and still get from every point to every other point. The theorem, however, asserts nothing about the distance one has to travel from  $x$  to  $y$  in the one-way system as compared to the distance between  $x$  and  $y$  in the two-way system. Actually, the comparison may be quite discouraging: if  $G$  is a cycle of length  $k$  then for each strongly connected orientation  $H$  of  $G$ , there are vertices  $x$  and  $y$  such that  $x$  and  $y$  are adjacent in  $G$  but it takes  $k-1$  edges to get from  $x$  to  $y$  in  $H$ .

J. A. Bondy and U. S. R. Murty proposed to study quantitative variations on Robbins' theorem. In particular, they conjectured the existence of a function  $f$  such that every bridgeless graph of diameter  $d$  admits an orientation of diameter  $f(d)$ . We shall prove their conjecture as a corollary to a rather general theorem. The theorem asserts that every undirected graph  $G$  admits an orientation  $H$  with the following property: if an edge  $uv$  belongs to a cycle of length  $k$  in  $G$  then  $uv$  or  $vu$  belongs to a cycle of length at most

$$(k-2)2^{\lfloor (k-1)/2 \rfloor} + 2$$

in  $H$ . It is an easy exercise to prove that, in a bridgeless graph  $G$  of diameter  $d$ , every edge belongs to a cycle of length at most  $2d+1$ .

Thus our theorem implies that the conjecture is true; in fact, it implies that

$$f(d) \leq d((2d-1)2^d + 1) .$$

This bound may be drastically improved. Indeed, we shall prove that every bridgeless graph of radius  $r$  admits an orientation of radius at most  $r^2+r$  (and this bound is best possible). It follows immediately that

$$f(d) \leq 2d^2 + 2d .$$

On the other hand we shall show that

$$f(d) \geq \frac{1}{2} d^3 - d$$

and we shall describe graphs of diameter  $d$  and arbitrary high (possibly infinite) connectivity such that every orientation has diameter at least

$$\frac{1}{4} d^2 + d .$$

Thus the order of growth of  $f$  is established; however, to find the exact values of  $f$  seems to be difficult. As a first step in this direction, we show that  $f(2) = 6$ ; the Petersen graph provides the lower bound. Finally, we shall turn our attention to the general problem of finding, for an undirected graph  $G$ , its orientation with a minimum diameter (resp. radius). We shall show that this problem is very difficult: in a sense, it is as difficult as the problem of deciding whether  $G$  has a hamiltonian cycle or the problem of finding the chromatic number of  $G$ .

In general, we follow the standard graph-theoretical notation and terminology, see Berge [2] or Harary [7]. In an undirected (resp.

directed) graph  $G$ , the distance  $\text{dist}(u,v;G)$  from  $u$  to  $v$  is the number of edges in a shortest path (resp. directed path) from  $u$  to  $v$ . Note that for an undirected graph  $G$ , the function  $\text{dist}(u,v;G)$  is a metric whereas for directed graphs  $G$ , we often have  $\text{dist}(u,v;G) \neq \text{dist}(v,u;G)$ . Unlike Moon [11], we postulate  $\text{dist}(u,u;G) = 0$ . The diameter of a graph  $G$  is the longest distance in  $G$ ; the radius of  $G$  is

$$\min_u \max_v (\max\{\text{dist}(u,v;G), \text{dist}(v,u;G)\}) .$$

Thus the diameter of  $G$  is at least the radius and at most twice the radius of  $G$ . Note that the diameter and the radius are defined only for connected undirected graphs and for strongly connected directed graphs. Also, if there is no finite upper bound on the distances in  $G$  then the diameter and radius are undefined.

1. From Cycles to Directed Cycles.

In the theorem below, we set

$$h(k) = (k-2)2^{\lfloor (k-1)/2 \rfloor} + 2$$

for every integer  $k$  such that  $k \geq 3$ .

THEOREM 1. Every graph  $G$  admits an orientation  $H$  with the following property: if an edge  $uv$  belongs to a cycle of length  $k$  in  $G$  then  $uv$  or  $vu$  belongs to a directed cycle of length at most  $h(k)$  in  $H$ .

PROOF. Let  $H_3$  be a maximal directed graph such that  $H_3$  is an orientation of some **subgraph** of  $G$  and such that every edge of  $H_3$  is in a directed cycle of length three. (If  $G$  is infinite then  $H_3$  exists by **Zorn's lemma**.) When  $H_{i-1}$  has been chosen for some  $i$ , let  $H_i$  be a maximal directed graph such that  $H_{i-1} \subset H_i$  and  $H_i$  is an orientation of some **subgraph** of  $G$  and **every** edge of  $H_i$  is in a directed cycle of length at most  $i$ . The graph

$$H = \bigcup_{i=3}^{\infty} H_i$$

is not necessarily an orientation of  $G$ : the bridges of  $G$  do not belong to  $H$ . However, we shall prove that  $H$  has the other desired property: if an edge  $uv$  belongs to a cycle of length  $k$  in  $G$  then  $uv$  or  $vu$  belongs to a directed cycle of length at most  $h(k)$  in  $H$ . Clearly, that is all we need: the edges belonging to no cycles of  $G$  (that is, the bridges of  $G$ ) may be directed quite arbitrarily.

Let us consider a cycle  $u_1, u_2, \dots, u_k, u_1$  in  $G$  such that, for some  $n$ , neither  $u_{1k}$  nor  $u_k u_1$  belongs to  $H_n$ ; all we have to prove is that  $n < h(k)$ .

For each  $i$  with  $3 < i \leq n$ , let  $x_i$  (resp.  $y_i$ ) denote the number of those directed edges  $u_{j+1}u_j$  (resp.  $u_j u_{j+1}$ ) that belong to  $H_i$  but not to  $H_{i-1}$ . For each  $m = 3, 4, \dots, n$ , we shall prove that

$$\sum_{i=3}^m (i-2)x_i \geq m+1-k \quad (1)$$

For this purpose, consider the graph  $H_m^*$  obtained from  $H_m$  by adding the directed edge  $u_k u_1$  and all the directed edges  $u_j u_{j+1}$  such that

$u_{j+1}u_j \notin H_m^*$ . Clearly, all the new edges of  $H_m^*$  lie on a directed cycle of length at most

$$1 + \sum_{j=1}^{k-1} \text{dist}(u_j, u_{j+1}; H_m^*) = \left( k - \sum_{i=3}^m x_i \right) + \sum_{i=3}^m (i-1)x_i .$$

By the maximality of  $H_m$ , this number is at least  $m+1$  and so (1) follows.

Next, define  $m(0) = 2$  and, for every positive integer  $t$ ,  $m(t) = (k-2)2^{t-1} + 2$ . For each  $t$  such that  $m(t) < n$ , we shall prove that

$$\sum_{i=3}^{m(t)} x_i \geq t . \quad (2)$$

Let  $t$  be the smallest nonnegative integer such that  $m(t) \leq n$  and such that (2) fails. Trivially,  $t \geq 1$ ; by the minimality of  $t$ , we have

$$t-1 \leq \sum_{i=3}^{m(t-1)} x_i \leq \sum_{i=3}^{m(t)} x_i \leq t-1$$

and so  $x_i = 0$  for  $m(t-1) < i \leq m(t)$ . Consequently,

$$\begin{aligned} \sum_{i=3}^{m(t)} (i-2)x_i &= \sum_{i=3}^{m(t-1)} (i-2)x_i \\ &\leq \sum_{s=1}^{t-1} (m(s)-2) \sum_{i=m(s-1)+1}^{m(s)} x_i = \sum_{s=1}^{t-1} (m(s) - m(s-1)) \sum_{i=m(s-1)+1}^{m(t-1)} x_i \\ &\leq \sum_{s=1}^{t-1} (m(s) - m(s-1))(t-s) = \sum_{s=1}^{t-1} m(s) - (t-1)m(0) \\ &= m(t) - k , \end{aligned}$$

contradicting (1). The same argument shows that, for each  $t$  such that  $m(t) \leq n$ , we have

$$\sum_{i=3}^{m(t)} y_i \geq t .$$

Now, we cannot have  $n \geq h(k) = m(\lfloor (k+1)/2 \rfloor)$  : indeed, this would imply

$$\sum_{i=3}^n x_i + \sum_{i=3}^n y_i \geq 2 \left\lfloor \frac{k+1}{2} \right\rfloor \geq k$$

which is clearly a contradiction.

COROLLARY 1. Let  $G$  be a graph such that every edge of  $G$  belongs to a cycle of length at most  $k$ . Then there is an orientation  $H$  of  $G$  such that

$$\text{dist}(u,v;H) \leq (h(k)-1)\text{dist}(u,v;G)$$

for every two vertices  $u$  and  $v$ .

A particular instance of Corollary 1 (to be used in Section 3) asserts the following: if every edge of  $G$  belongs to a triangle then there is an orientation  $H$  of  $G$  such that

$$\text{dist}(u,v;H) \leq 3 \cdot \text{dist}(u,v;G) .$$

for every two vertices  $u$  and  $v$ .

Note that  $h(3) = 4$  and  $h(4) = 6$ . If  $h'$  is any function such that Theorem 1 holds with  $h'$  instead of  $h$ , then we must have  $h'(3) \geq 4$  (as demonstrated by a wheel with an odd number of spokes) and  $h'(4) \geq 6$  (as demonstrated by the pentagonal prism) so for  $k = 3, 4$  the result of Theorem 1 is best possible.

However, we do not know if  $h'$  can be chosen to be a polynomial or even a linear function.

## 2. From Radius to Directed Radius.

THEOREM 2. Every bridgeless graph of radius  $r$  admits an orientation of radius at most  $r^2+r$ .

PROOF. We shall find it useful to work with orientations of multi-graphs: in such orientations, the multiple edges may be directed both ways (whereas the single edges must be directed only one way). By induction on  $r$ , we shall prove the following statement: "if  $G$  is a bridgeless multigraph and if  $u$  is a vertex of  $G$  such that  $\text{dist}(u,v;G) \leq r$  for every vertex  $v$  then there is an orientation  $H$  of  $G$  such that  $\text{dist}(u,v;H) \leq r^2+r$  and  $\text{dist}(v,u;H) \leq r^2+r$  for every vertex  $v$ ".

For every neighbor  $v$  of  $u$ , the edge  $uv$  is contained in some cycle; let  $k(v)$  denote the length of a shortest such cycle. It is important to note that

$$k(v) \leq 2r+1 \quad \text{for every } v; \quad (\text{3})$$

the proof of this fact is left to the reader. An orientation  $A$  of some subgraph of  $G$  will be called admissible if there is a set  $S$  of neighbors of  $u$ , together with a directed cycle  $C_v$  for each  $v \in S$ , such that

- (i) each  $C_v$  has length  $k(v)$  and contains either the edge  $uv$  or the edge  $vu$ ,

(ii)  $A$  is the union of all these cycles  $C_v$  ( $v \in S$ ).

Note that by (3) and this definition, we have

$$\text{dist}(u, w; A) \leq 2r \quad \text{and} \quad \text{dist}(w, u; A) \leq 2r \quad (4)$$

for every vertex  $w$  of  $A$ . Furthermore, we shall prove that

$$\begin{aligned} &\text{every maximal admissible graph} \\ &\text{contains all the neighbors of } u. \end{aligned} \quad (5)$$

Assume the contrary, so that  $w \notin A$  for some maximal admissible graph  $A$  and for some neighbor  $w$  of  $u$ . There is a cycle

$w_1, w_2, \dots, w_k, w_1$  in  $G$  such that  $w_1 = u$ ,  $w_2 = w$  and  $k = k(w)$ .

If none of the vertices  $w_2, w_3, \dots, w_k$  belongs to  $A$  then adding the

directed circuit  $w_1 \rightarrow w_2 \rightarrow \dots \rightarrow w_k \rightarrow w_1$  to  $A$  we obtain a larger admissible graph: a contradiction. Thus we may assume that at least

one of the vertices  $w_2, w_3, \dots, w_k$  belongs to  $A$ ; let  $w_i$  be such a vertex with the smallest subscript. Since  $w_i \in A$ , there is some  $v \in S$

such that  $w_i \in C_v$ . Writing  $u = v_1 \rightarrow v_2 \rightarrow \dots \rightarrow v_m \rightarrow v_1$  for  $C_v$ ,

we have  $m = k(v)$  and either  $v = v_2$  or  $v = v_m$ ; there is no loss of

generality in assuming that  $v = v_2$ . We also have  $w_i = v_j$  for some  $j$ .

Now, we shall distinguish between two cases.

Case 1.  $w_k = v$ . In this case, define  $C_w$  to be the directed cycle

$$u \rightarrow v_2 \rightarrow v_3 \rightarrow \dots \rightarrow v_j \rightarrow w_{i-1} \rightarrow w_{i-2} \rightarrow \dots \rightarrow w_2 \rightarrow u$$

and note that  $C_w$  has length  $k(w)$ . (Indeed, if  $C_w$  had more than

$k(w)$  edges then the path  $u, v, v_3, \dots, v_j$  would be longer than the path

$u, v, w_{k-1}, \dots, w_i$ . In that case, the closed walk

$u, v, w_{k-1}, \dots, v_{j+1}, \dots, v_m, u$  would produce a cycle in  $G$  of length

less than  $k(v)$  and yet containing the edge  $uv$  : a contradiction.)

Adding  $C_w$  to  $A$  we obtain a larger admissible graph: a contradiction.

Case 2.  $w_k \neq v$ . In this case, define  $C_w$  to be the directed cycle

$$u \rightarrow w_2 \rightarrow w_3 \rightarrow \dots \rightarrow w_i \rightarrow v_{j+1} \rightarrow v_{j+2} \rightarrow \dots \rightarrow v_m \rightarrow u$$

and note that  $C_w$  has length  $k(w)$ . (Indeed, if  $C_w$  had more than  $k(w)$  edges then the path  $v_j, v_{j+1}, \dots, v_m, u$  would be longer than the path  $w_i, w_{i+1}, \dots, w_k, u$ . In that case, the closed walk  $u, v, v_j, w_i, w_{i+1}, \dots, w_k, u$  would produce a cycle in  $G$  of length less than  $k(v)$  and yet containing the edge  $uv$  : a contradiction.)

Adding  $C_w$  to  $A$  we obtain a larger admissible graph: a contradiction.

Now, (5) is proved and the rest is fairly straightforward. Consider a maximal admissible graph  $A$ . (If  $G$  is infinite then the existence of  $A$  follows by Zorn's lemma.) In  $G$ , contract all the vertices of  $A$  into a new vertex  $u^*$  (this may create new multiple edges) and call the resulting graph  $G^*$ . Note that  $G^*$  is bridgeless and that by (5), we have

$$\text{dist}(u^*, v; G) \leq r-1$$

for every vertex  $v$  of  $G^*$ . By the induction hypothesis, there is an orientation  $H^*$  of  $G^*$  such that

$$\text{dist}(u^*, v; H^*) \leq r^2 - r \quad \text{and} \quad \text{dist}(v, u^*; H^*) \leq r^2 - r \quad (6)$$

for every vertex  $v$  of  $G^*$ . We may think of  $H^*$  as an orientation of some subgraph of  $G$ . Continuing this orientation with  $A$  (and directing all the remaining edges of  $G$  arbitrarily) we obtain an orientation  $H$  of  $G$ . By (4) and (6), we have

$$\text{dist}(u,v;H) \leq r^2+r \quad \text{and} \quad \text{dist}(v,u;H) \leq r^2+r$$

for every vertex  $v$  of  $G$ .

THEOREM 3. For every positive integer  $r$  there is a bridgeless graph  $G_r$  of radius  $r$  such that every orientation of  $G_r$  has radius  $r^2+r$ .

PROOF. We shall construct a certain sequence  $H_1, H_2, \dots$  of rooted graphs.  $H_2$  is simply a triangle with one of its vertices designated as the root. To construct  $H_r$ , take a cycle  $u_0, u_1, \dots, u_{2r}, u_0$  and two disjoint copies of  $H_{r-1}$ . Then identify the root of the first (resp. second) copy of  $H_{r-1}$  with  $u_1$  (resp.  $u_{2r}$ ). The resulting graph, rooted at  $u_0$ , is  $H_r$ . Finally,  $G_r$  is obtained by taking two disjoint copies of  $H_r$  and identifying their roots. The graph  $G_3$  is shown in Figure 1; we leave it to the reader to verify that  $G_r$  has the desired property.




Figure 1<sup>i</sup>

**REMARK.** The graphs  $G_r$  constructed above are so easy to study because of their simple structure and numerous cutpoints. We do not know if there are undirected graphs  $G$  of arbitrarily high connectivity and radius  $r$  such that every orientation of  $G$  has radius at least  $r^2+r$ . Nevertheless, we can construct undirected graphs  $G$  of arbitrarily high connectivity and radius  $r$  such that every orientation of  $G$  has radius at least  $r^2/2+r$ . These are the graphs  $G_{2r,k}$  constructed in Theorem 4.

### 3. From Diameter to Directed Diameter.

For each positive integer  $d$ , let  $f(d)$  be the smallest integer such that every bridgeless graph of diameter  $d$  admits an orientation of diameter at most  $f(d)$ . By Theorem 2, we have  $f(d) \leq 2d^2+2d$ . On the other hand, the reader may verify that each of the graphs  $G_r$  of Theorem 3 has diameter  $d = 2r$  and that every strongly connected orientation of  $G_r$  has diameter  $2(r^2+r) = d^2/2+d$ . Let  $G'_r$  denote the graph obtained from  $H_r$  and  $H_{r+1}$  (of Theorem 3) by identifying their roots. Then  $G'_r$  has diameter  $d = 2r+1$  and every strongly connected orientation of  $G'_r$  has diameter  $r^2+r+(r+1)^2+(r+1) = \frac{1}{2}d^2+d+\frac{1}{2}$ . Hence  $f(d) \geq \frac{1}{2}d^2+d$  for all  $d \geq 2$ .

**THEOREM 4.** For each pair  $d, k$ , where  $d$  is a positive integer and  $k$  is a finite or infinite cardinal, there is a  $k$ -connected undirected graph  $G_{d,k}$  of diameter  $d$  such that every orientation of  $G_{d,k}$  has diameter at least  $\frac{1}{4}d^2+d$ .

PROOF. Begin with disjoint sets of vertices  $S_1, S_2, \dots, S_m$  such that  $S_1$  and  $S_m$  have cardinality 1, each of  $S_i$  ( $2 \leq i \leq m-1$ ) has cardinality  $k$ , and

$$m = \begin{cases} 1 + (d+1)^2/2 & (d \text{ odd}) , \\ 1 + d(d+2)/2 & (d \text{ even}) . \end{cases}$$

Then, for each  $i = 1, 2, \dots, m-1$ , join every vertex from  $S_i$  to every vertex from  $S_{i+1}$ . The resulting graph is  $k$ -connected; by adding as few as  $d$  edges, we shall bring its diameter down to  $d$ . To do so, we shall first define

$$i(j) = \begin{cases} 1 + (j+1)j & \text{for } 0 \leq j \leq d/2 , \\ m - (d-j+1)(d-j) & \text{for } d/2 \leq j \leq d . \end{cases}$$

Note that  $i(0) = 1 < i(1) < \dots < i(d-1) = m-2$  and that the sequence of differences  $i(j+1) - i(j)$  is  $2, 4, 6, \dots, 6, 4, 2$ . For each

$j = 0, 1, \dots, d$  choose a vertex  $u_j \in S_{i(j)}$ . For each  $j = 0, 1, 2, \dots, d-1$  join  $u_j$  to  $u_{j+1}$ . Call the resulting graph  $G_{d,k}$ . The graph  $G_{4,2}$  is shown in Figure 2. It is easy to see that  $G_{d,k}$  has diameter  $d$ .

Now consider any strongly connected orientation of  $G_{d,k}$ . Let  $P_1$  (resp.  $P_2$ ) denote a shortest directed path from  $u_0$  to  $u_d$  (resp. from  $u_d$  to  $u_0$ ).

Let  $l_s$  denote the length of  $P_s$  for  $s = 1, 2$ .

For each  $j$ ,  $0 \leq j \leq d-1$ , at least one of the paths  $P_s$ ,

$1 \leq s \leq 2$ , contains a subpath from  $S_{i(j)}$  to  $S_{i(j+1)}$  avoiding the

edge between  $u_j$  and  $u_{j+1}$ , so  $l_1 + l_2 \geq \sum_{j=0}^{d-1} (i(j+1) - i(j) + 1) =$

$m-1+d$ . Hence the diameter is at least  $\frac{1}{2}(m-1+d) \geq \frac{1}{4}d^2 + d$ .




Figure 2

In the rest of this section, we shall prove  $f(2) = 6$ .

THEOREM 5. Every bridgeless graph of diameter two admits an orientation of diameter at most six.

PROOF. Let  $G$  be a bridgeless graph of diameter two. We may assume that some edge  $uv$  of  $G$  is contained in no triangle (otherwise the desired conclusion follows from Corollary 1). Let  $A$  (resp.  $B$ ) denote the set of all the neighbors of  $u$  (resp.  $v$ ) other than  $v$  (resp.  $u$ ). Furthermore, let  $A_1$  (resp.  $B_1$ ) denote the set of all the vertices in  $A$  (resp.  $B$ ) that have no neighbors in  $B$  (resp.  $A$ ). Set  $A_2 = A - A_1$ ,  $B_2 = B - B_1$  and denote by  $C$  the set of all the vertices not in  $\{u, v\} \cup A \cup B$ . The reader may easily verify that the orientation of  $G$ , described simply by  $u \rightarrow v \rightarrow B \rightarrow C \rightarrow A \rightarrow u$  and  $B_1 \rightarrow B_2 \rightarrow A_2 \rightarrow A_1$ , has diameter at most six. (Here  $X \rightarrow Y$  means that every edge joining a vertex  $x$  of  $X$  with a vertex  $y$  of  $Y$  is directed from  $x$  to  $y$ .)

Next, we shall prove that Theorem 5 is best possible. A part of our argument is of independent interest; therefore we shall state it on its own.

LEMMA. Every strongly connected orientation of the Petersen graph contains a directed cycle of length five.

PROOF. Let  $H$  be a strongly connected orientation of the Petersen graph; assume that  $H$  contains no directed cycle of length five. Since  $H$  is strongly connected, it contains some directed cycle; furthermore, the shortest directed cycle has no diagonals. In the Petersen graph, there are no cycles of length seven and every cycle of length greater than seven has a diagonal. Hence we may assume that  $H$  contains a directed cycle of length six. In the Petersen graph, every two cycles of length six are equivalent under some automorphism; hence we may assume that  $H$  contains the directed cycle  $1 \ 4 \ 2 \ 4 \ 3 \ 4 \ 4 \ 4 \ 5 \ 4 \ 6 \ 4 \ 1$  shown in Figure 3.




Figure 3

At least one edge of  $\Pi$  enters 0 and at least one edge of  $\Pi$  leaves 0 ; without loss of generality, we may assume  $7 \rightarrow 0$  and  $8 \rightarrow 5$ , creating the directed cycle  $5 \rightarrow 0 \rightarrow 1 \rightarrow 2 \rightarrow 3 \rightarrow 4 \rightarrow 5$  (otherwise  $8 \rightarrow 2$  forces

Similarly,

we must have  $3 \rightarrow 7$  (otherwise  $7 \rightarrow 3$  forces  $6 \rightarrow 7$  creating the directed cycle  $3 \rightarrow 4 \rightarrow 5 \rightarrow 6 \rightarrow 7 \rightarrow 3$ ). But then  $3 \rightarrow 7 \rightarrow 0 \rightarrow 8 \rightarrow 2 \rightarrow 3$  is a directed cycle of length five: a contradiction.

(Let us digress in order to mention a problem suggested by the lemma. Which bridgeless graphs  $G$  have the property that every strong orientation of  $G$  contains a directed cycle whose length equals the girth of  $G$ ? The Petersen graph has this property and so does every bridgeless graph of radius one.)

THEOREM 6. Every orientation of the Petersen graph has diameter at least six.

PROOF. Let us assume that some orientation  $H$  of the Petersen graph has diameter at most five. By the lemma,  $H$  contains a directed cycle of length five. In the Petersen graph, every two cycles of length five are equivalent under some automorphism; hence we may assume that  $H$  contains the cycle  $1 \rightarrow 2 \rightarrow 3 \rightarrow 4 \rightarrow 5 \rightarrow 1$  shown in Figure 4.




Figure 4

Consider the "cross edges"  $16, 27, 38, 49, 50$ ; each of them may be directed towards the pentagon or away from it.

Case 1. Three consecutive cross edges are directed in the same sense. Without loss of generality, we may assume that  $0 \rightarrow 5$ ,  $6 \rightarrow 1$  and  $7 \rightarrow 2$ . Here, the contradiction is immediate:  $\text{dist}(5,7;H) \geq 6$ .

Case 2. No three consecutive edges are directed in the same sense. We may assume  $8 \rightarrow 3$  and  $9 \rightarrow 4$ , forcing  $5 \rightarrow 0$ ,  $2 \rightarrow 7$  and, in turn,  $6 \rightarrow 1$ . Now,  $\text{dist}(3,6;H) \leq 5$  forces  $0 \rightarrow 8 \rightarrow 6$  and  $\text{dist}(3,9;H) < 5$  forces  $0 \rightarrow 7 \rightarrow 9$ . But then  $\text{dist}(1,8;H) \geq 6$ , a contradiction.

We do not know any other bridgeless graph of diameter 2 which can play the role of the Petersen graph in Theorem 6. Perhaps the Moore graph of diameter two and degree seven, constructed by Hoffman and Singleton [8], is another example.

#### 4. Finding Optimum Orientations.

Given an undirected graph  $G$ , let us ask for its orientation with smallest possible diameter (resp. radius). Clearly, such an orientation can be found in a finite time: the diameter and the radius of a directed graph on  $n$  vertices can be found in  $O(n^2)$  steps [6] and an undirected graph with  $m$  edges has  $2^m$  distinct orientations. In the spirit of Edmonds [5], we shall ask for a "better-than-finite" algorithm finding optimum orientations; more specifically, we shall ask for such an algorithm terminating within  $p(n)$  steps for some fixed polynomial  $p$ . Our results are rather discouraging; "they suggest that no such an algorithm exists.

The key notion is that of an "NP-hard" problem. A certain class of problems is called NP. This class is very wide: it consists of all the problems for which the correctness of a proposed solution may be checked in a polynomial time (relative to the size of the problem). That is, NP consists of all the problems with "good characterizations" (this notion is due to Edmonds [4]). For example, the problem "Is a graph  $G$   $k$ -colorable?" belongs to NP. Now, let  $X$  be a problem with the following property: if  $X$  can be solved in a polynomial time then every problem in NP can be solved in a polynomial time. Such a problem is called NP-hard. (To many, it seems unlikely that every problem in NP can be solved in a polynomial time; such a belief implies that no NP-hard problem can be solved in a polynomial time.) In a pioneering paper [3], Cook proved that it is NP-hard to find the largest clique in a graph. Since then, many other people have shown many other problems to be NP-hard; as a rule, this is done by "reducing"

the problem of finding the largest clique in a graph (or another problem which already has been shown to be NP-hard) into the problem in question. (For more information on the subject, the reader is referred to [1].)

In particular, Lovász [10] has shown that it is NP-hard to decide if a hypergraph is 2-colorable; it is implicit in his proof that the same problem remains NP-hard even when the input is restricted to hypergraphs of rank three. The relevant definitions may be found in Berge's monograph [2]; for the sake of completeness, we shall repeat them here. A hypergraph is an ordered pair  $H = (V, E)$  such that  $V$  is a set and such that  $E$  is a family of subsets of  $V$ . The elements of  $V$  are called the vertices of  $H$ , the elements of  $E$  are called the edges of  $H$ . The number of vertices of  $H$  is called the order of  $H$ , the cardinality of the largest edge of  $H$  is called the rank of  $H$ . A hypergraph is called 2-colorable if its vertices can be colored red and blue in such a way that every edge includes at least one vertex of each color.

**THEOREM 7.** Given a hypergraph  $H$  of rank three and order  $n$ , we can construct in  $O(n^6)$  steps a graph  $G$  with the following property:  $G$  admits an orientation of diameter two if and only if  $H$  is 2-colorable.

**PROOF.** Let  $k$  be the integer satisfying  $10 \leq k \leq 13$  and  $n+k \equiv 2 \pmod{4}$ . We shall find it convenient to work with the hypergraph  $H_0$ , obtained from  $H$  by adding  $k$  new vertices  $v_1, v_2, \dots, v_k$  and, if  $H$  has an even number of edges, adding new edge  $\{v_1, v_2\}$ . Note that  $H_0$  has an odd number of edges. To

construct  $G$ , take disjoint sets  $P$  and  $Q$  such that the elements of  $P$  (resp.  $Q$ ) are in a one-to-one correspondence with the vertices (resp. the edges) of  $H_0$ ; for simplicity, we shall label each element of  $P$  (resp.  $Q$ ) by the corresponding vertex (resp. edge) of  $H_0$ . Join by an edge every two vertices in  $P$  and every two vertices in  $Q$ ; join a vertex  $v \in P$  to a vertex  $e \in Q$  if and only if  $v \in e$  in  $H_0$ . Then add four vertices  $w_1, w_2, w_3, w_4$  and join each of them to all the vertices in  $P \cup Q$ . Finally, add a new vertex  $x$  and join it to all the vertices in  $P$ . We shall show that the resulting graph  $G$  has the desired property. (Note that the number of edges of  $G$  may be of the order  $\binom{6}{n}$ .)

Firstly, assume that  $G$  admits an orientation  $G^*$  of diameter two. Color a vertex  $u$  of  $H$  blue (resp. red) if in  $G^*$ , we have  $x \rightarrow u$  (resp.  $u \rightarrow x$ ). Since  $\text{dist}(x, e; G^*) = 2$  (resp.  $\text{dist}(e, x; G^*) = 2$ ) for every  $e \in Q$ , every edge of  $H$  includes at least one blue (resp. red) vertex. Thus  $H$  is 2-colorable.

Secondly, assume that  $H$  is 2-colorable. Then  $H_0$  admits a 2-coloration such that the number of blue (and red) vertices is odd and at least five; this 2-coloration induces a partition  $P = P_1 \cup P_2$ . We are going to describe an orientation  $G^*$  of  $G$ ; before doing so, let us digress a little. By a cyclic tournament of order  $2k+1$ , we shall mean the tournament with vertices  $u_1, u_2, \dots, u_{2k+1}$  such that  $u_1 \rightarrow u_{i+j}$  for every  $j = 1, 2, \dots, k$  (arithmetic modulo  $2k+1$ ). The parity partition of such a tournament is the partition  $A \cup B$  defined by  $A = \{u_1, u_3, \dots, u_{2k+1}\}$  and  $B = \{u_2, u_4, \dots, u_{2k}\}$ . If  $k \geq 2$  then the parity partition has the following nice properties:

(i) if  $u \in A$  then there are  $v_1, v_2 \in A$  such that  $v_1 \rightarrow u \rightarrow v_2$ ,

(ii) if  $u \in B$  then there are  $v_1, v_2 \in B$  such that  $v_1 \rightarrow u \rightarrow v_2$ .

Now, the orientation  $G^*$  of  $G$  may be described as follows. For each of the three sets  $P_1, P_2, Q$ , direct the edges of the complete graph induced by that set so as to obtain a cyclic tournament. Let  $A_i \cup B_i$  be the parity partition of  $P_i$  ( $i = 1, 2$ ) and let  $A \cup B$  be the parity partition of  $Q$ . Direct

$$\begin{aligned}
 x &\rightarrow P_1 \rightarrow P_2 \rightarrow x, \\
 P_1 &\rightarrow Q \rightarrow P_2, \\
 A_1 \cup A_2 &\rightarrow w_1 \rightarrow A, \\
 B &\rightarrow w_1 \rightarrow B_1 \cup B_2, \\
 A_1 \cup A_2 &\rightarrow w_2 \rightarrow B, \\
 A &\rightarrow w_2 \rightarrow B_1 \cup B_2, \\
 B_1 \cup B_2 &\rightarrow w_3 \rightarrow A, \\
 B &\rightarrow w_3 \rightarrow A_1 \cup A_2, \\
 B_1 \cup B_2 &\rightarrow w_4 \rightarrow B, \\
 A &\rightarrow w_4 \rightarrow A_1 \cup A_2.
 \end{aligned}$$

We leave it to the reader to verify that  $G^*$  has diameter two.

COROLLARY 2. It is NP-hard to decide whether an undirected graph admits an orientation of diameter two.

THEOREM 8. Given a hypergraph  $H$  of rank three and order  $n$ , we can construct in  $O(n^6)$  steps a graph  $G$  with the following property:  $G$  admits an orientation of radius two if and only if  $H$  is 2-colorable.

PROOF. Take disjoint sets  $P$  and  $Q$  such that the elements of  $P$  (resp.  $Q$ ) are in a one-to-one correspondence with the vertices (resp. the edges) of  $H$ . Join by an edge every two vertices in  $P$ ; join a vertex  $v \in P$  to a vertex  $e \in Q$  if and only if  $v \in e$  in  $H$ . Then add a new vertex  $x$  and join it to all the vertices in  $P$ ; call the resulting graph  $G_0$ . To construct  $G$ , take two disjoint copies of  $G_0$  and identify their vertices  $x$ . We leave it to the reader to verify that  $G$  has the desired property.

COROLLARY 3. It is NP-hard to decide whether an undirected graph admits an orientation of radius two.

REMARK. Easy modifications of our constructions show that (i) for every  $k$  with  $k \geq 4$ , it is NP-hard to decide whether an undirected graph admits an orientation with diameter at most  $k$ , (ii) for every  $k$  with  $k \geq 4$ , it is NP-hard to decide whether an undirected graph  $G$  admits an orientation with radius at most  $k$ . To prove (i), take the graph  $G$  as constructed in Theorem 7, add a cycle  $u_0, u_1, \dots, u_{k-2}, u_0$  and identify  $u_0$  with  $x$ . To prove (ii), take two disjoint copies of the graph  $G_0$  constructed in Theorem 8, add a tree  $T$  consisting of four paths of length  $k-2$  starting at the same vertex. Now take two of the four end vertices of  $T$  and identify them with the vertex  $x$  in one of the copies of  $G_0$ , then identify the remaining two end vertices of  $T$  with the vertex  $x$  in the other copy of  $G_0$ .

**REMARK.** Corollary 2 shows that in general, it is very hard to decide whether an undirected graph  $G$  admits an orientation of diameter two. However, if  $G$  has too few edges then the answer is always negative. More precisely, Katona and Szemerédi[9] proved that no undirected graph with  $n$  vertices and fewer than  $\frac{n}{2} \log_2 \frac{n}{2}$  edges admits an orientation of diameter two.

## References

- [1] A. V. Aho, J. E. Hopcroft and J. D. Ullman, The Design and Analysis of Computer Algorithms, Addison-Wesley, Reading, Mass., **1.974**, Chapter 10.
- [2] C. Berge, Graphs and Hypergraphs, North-Holland, Amsterdam, 1973.
- [3] S. A. Cook, "The complexity of theorem proving procedures," Proc. Third Annual ACM Symposium on Theory of Computing, 151-158.
- [4] J. Edmonds, "Minimum partition of a matroid into independent subsets," J. Res. Nat. Bur. Standards B69 (1965), **67-72**.
- [5] J. Edmonds, "Paths, trees, and flowers," Canad. J. Math. 17 (**1965**)' **449-467**.
- [6] R. W. Floyd, "Algorithm 97: shortest path," Comm. ACM 5 (1962), **345**.
- [7] F. Harary, Graph Theory, Addison-Wesley, Reading, Mass., 1969.
- [8] A. J. Hoffman and R. R. Singleton, "On Moore graphs with diameter 2 and 3," IBM J. Res. Develop. 4 (1960), 497-504.
- [9] G. Katona and E. Szemerédi, "On a problem of graph theory," Studia Sci. Math. Hungar. 2 (1967), **23-28**.
- [10] L. Lovász, "Coverings and colorings of hypergraphs," Proc. Fourth S.-E. Conf. Combinatorics, Graph Theory and Computing (F. Hoffman et al., eds.), Utilitas Mathematica Publishing Inc., Winnipeg, **1973**, **3-12**.
- [11] J. W. Moon, Topics on Tournaments, Holt, Rinehart and Winston, New York, **1968**; Section 12.
- [12] H. E. Robbins, "A theorem on graphs with an application to a problem of traffic control," Am. Math. Monthly 46 (1939), 281-283.